

## Acute toxic effects of cadmium in larvae of the green toad, *Pseudepidalea variabilis* (Pallas, 1769) (Amphibia: Anura)

Mert Gürkan, Ayşe Çetin, and Sibel Hayretdağ

Department of Biology, Zoology Section, Faculty of Arts and Sciences, Çanakkale Onsekiz Mart University, Turkey

Received in April 2014

CrossChecked in April 2014

Accepted in August 2014

The environmental impact of cadmium use and its accumulation in nature have increased to alarming levels. This study aimed to morphologically and histologically investigate the acute toxic effects of cadmium on green toad, *Pseudepidalea variabilis* (Pallas, 1769) larvae. Embryos were obtained from specimens collected in amplexus from nature and kept under laboratory conditions until stage 26, when they were exposed to cadmium (0, 1, 5, 10, 25, and 50  $\mu\text{g L}^{-1}$ ) for 96 h. The  $\text{LC}_{10}$ ,  $\text{LC}_{50}$ , and  $\text{LC}_{90}$  values of cadmium were calculated to be 26.98, 35.35, and 46.31  $\mu\text{g L}^{-1}$ , respectively. Our results showed that cadmium had a negative effect on the body size of *P. variabilis* larvae (over 1  $\mu\text{g L}^{-1}$ ). Histological examination detected a fusion of gill lamellae, liver haemorrhage, oedema in the abdominal cavity, and deformations of pronephric tubules (over 10  $\mu\text{g L}^{-1}$ ). Our findings suggest that the green toad was sensitive to the cadmium treatment, with  $\text{LC}_{50}$  values lower than those reported by other studies. Thus, this species could be considered a reliable indicator species of environmental stress in aquatic ecosystem.

**KEY WORDS:** *embryotoxicity; heavy metal; histopathology; lethal concentrations*

The decline in amphibian populations has reached striking dimensions in recent years (1, 2). There are several hypotheses about the possible reasons behind this phenomenon (3, 4). Some of these hypotheses include impact of urbanization due to rapid increase in settlements and industrialization, chemical pollution, habitat degradation/destruction, and diseases (5-8). Among these, the destruction of habitats and increased pollution significantly threaten the survival of amphibians, which spend their embryonic and larval stages exclusively in water and play a significant role in biomonitoring programmes (9-11). That is why amphibian larvae are among the preferred bioindicators for aquatic toxicology studies (12-16).

Cadmium is present in the environment naturally; however, it is not an essential element for organisms (17). Increases in the amount of cadmium in nature might be either due to natural factors, like the ablation

of rocks resulting from erosion and rain, or anthropogenic factors (18). In Turkey, for instance, one study (19) found the mean cadmium concentration in surface waters to be 110  $\mu\text{g L}^{-1}$ .

The half-life of cadmium in organisms is 20 years (20). It cannot be excreted from the body and accumulates in the organism and throughout the food chain. Its biomagnification poses threats for human health as well as the ecosystem as a whole. Therefore, cadmium has recently become the subject of many investigations (16, 21-25). The toxic effects of cadmium investigated in freshwater fish include development abnormalities detected in many teleost species (26-28). Cadmium led to larval teratogenic and developmental abnormalities (29), to micronucleus induction at environmental levels of 2, 10, and 30  $\mu\text{g L}^{-1}$ , and to increased concentration-dependent quantities of micronucleus after contamination with 10 and 30  $\mu\text{g L}^{-1}$  in *Xenopus* larvae (30). In addition,

cadmium exposure caused a decline in the survival numbers of *Bufo americanus* (13), developmental delays in tadpoles of *Bufo raddei* (31), and toxic acute effects on early-life stages in *Duttaphrynus melanostictus* at levels of 0.2 mg L<sup>-1</sup> (23). On the contrary, cadmium stimulated growth and development in amphibian larvae at concentrations from 0.25 to 5 µg L<sup>-1</sup> (32).

Known as the green toad, *Pseudepidalea variabilis* is a common amphibian species in Turkey. This terrestrial and nocturnal species is water-dependent in the breeding season, when it is known to stay in water for long periods of time (33). This species generally inhabits slow-flowing and stagnant waters, seasonal ponds, and shallow pits filled with water for egg-laying (34). Its aquatic habitats are prone to increased heavy metal concentrations due to agricultural and industrial activities. To our knowledge, until now no related study on the acute toxic effects of cadmium on *P. variabilis* larvae has been conducted.

This study aimed to determine the lethal concentrations (LC<sub>10</sub>, LC<sub>50</sub>, and LC<sub>90</sub>) of cadmium and the acute toxic effects on the tissues and organs of *P. variabilis* larvae after a 96-h acute exposure.

## MATERIALS AND METHODS

### *Animals and treatment*

The green toad, *Pseudepidalea variabilis* (Pallas, 1769), is included in the “data deficient” (DD) category of the IUCN Red List (35). Tadpoles were obtained in the laboratory from 4 adult (2 ♂♂, 2 ♀♀) *P. variabilis* caught in amplexus in March 2010, in a seasonal pond in Çanakkale, north-western Turkey. *P. variabilis* specimens were transferred to the laboratory and kept in polypropylene containers until ovulation. After ovulation, embryos were transferred to 30×30×25 cm glass aquaria. After hatching, the larvae were fed with fish feed of plant origin (100 g/ carbohydrate: 30.16 g; sugars: 7.33 g; dietary fibre: 9.3 g; fat: 19.94 g; protein: 36.49 g; energy: 450 kcal) and boiled lettuce leaves until stage 26 (36).

Cadmium chloride (CdCl<sub>2</sub>) was supplied from Sigma-Aldrich Chemical Company Inc. (St. Louis, MO, USA). A stock solution of 100 µg L<sup>-1</sup> was prepared by dissolving CdCl<sub>2</sub> powder in distilled water. Larvae at stage 26 were exposed to concentrations of 1, 5, 10, 25, and 50 µg L<sup>-1</sup> of cadmium. The dosing

solutions were prepared by appropriate dilutions of the stock solution.

Twenty randomly selected *P. variabilis* larvae were exposed to control (water) and each cadmium concentration, in acute (96 h) exposure experiments, in a static bioassay test system. Aquaria were filled with 2 L of water or cadmium solutions and during exposure, temperature, pH, and dissolved oxygen levels were measured daily with an Elmetron CO-401 meter (analysio GmbH, Greifswald, Germany). The treatment was carried out in a 14:10-h light-dark cycle and the number of dead tadpoles was recorded and faeces removed every 24 hours.

### *Morphology and histology*

Ten randomly selected larvae from the control and each treatment group were measured for total length, width, and tail length. Furthermore, wet weights of the larvae were determined (0 and 96 h). Wet weight (precision 0.001 g) and total length (precision 0.01 mm) were measured after euthanasia with a digital scale and digital calipers. Ten larvae survived after the 96 h of exposure were fixed in Bouin's solution for histological examination after macroscopic evaluation. Afterwards, these larvae were processed through alcohol, xylene, and paraffin series and then paraffin blocks were prepared. Serial cross sections 6-8 µm in thickness were obtained from these blocks, stained with hematoxylin and eosin (H&E), and histopathologically examined under a light microscope. Finally, the histological imaging of the preparations was carried out using a camera mounted on an Olympus BX51 light microscope (Japan) and analysed using the DP2-BSW software.

### *Statistical analysis*

Finney's Probit analysis method was used to calculate 96-h LC<sub>10</sub>, LC<sub>50</sub>, and LC<sub>90</sub> values of cadmium for *P. variabilis* larvae. Lethal concentrations were calculated at 95 % confidence intervals. Comparisons of total length, tail length, width and weight between control and treatment groups were done using one-way analysis of variance (ANOVA). Discriminant analysis was done in order to show the possible total differences of the concentrations tested. Statistical analyses were performed using SPSS software (ver. 16.0) and alpha was set at 0.05.

## RESULTS

### Acute toxicity

Mean water temperature was  $21.2 \pm 2$  °C, while pH and dissolved oxygen in aquaria were  $6.8 \pm 0.3$  and  $8.2 \pm 0.4$  mg L<sup>-1</sup>, respectively. No mortality was recorded in the control group and in the cadmium groups exposed to 1 and 5 µg L<sup>-1</sup>. The highest mortality was observed at 50 µg L<sup>-1</sup>, where 10 larvae died within the first 24 h and only one larva survived at the end of the exposure period. After 72 h, a dead larva was found in the 10 µg L<sup>-1</sup> group, whereas 8 larvae died in the 25 µg L<sup>-1</sup> group. Values of 96-h LC<sub>10</sub>, LC<sub>50</sub>, and LC<sub>90</sub> are presented in Table 1.

### Morphology and histology


Total length was the only parameter found to be significantly different between treatments ( $p < 0.05$ ). Comparisons of total length, width, tail length, and wet weight between control and treatment groups are presented in Table 2. Since only one larva survived the 50 µg L<sup>-1</sup> concentration to the end, this group was not included in the statistical analysis.

The canonic discriminant analysis was performed to assess differences in total body length among groups. It showed two functions, which explained 100 % of the total variance (Figure 1) and the  $p$ -value of function 1 was significant (Table 3).

No abnormalities were found in the histological examinations of larvae from the control group. Likewise, no significant histological findings were detected in the groups exposed to 1 and 5 µg L<sup>-1</sup> of cadmium. However, deformations and gill lamellar fusions were recorded in some larvae at 10 and 25 µg L<sup>-1</sup> (Figure 2).

Furthermore, effects such as deformations of pronephric tubules structure were prominent and clearly observed at increasing concentrations (Figure 3).

No histopathological alterations were encountered in the cross sections of the livers of larvae in the



**Figure 1** According to total length measurements, the total differences of exposure concentrations between groups

control group. Nevertheless, deformation and haemorrhage in liver were observed in most of the larvae exposed to 10 and 25 µg L<sup>-1</sup> of cadmium. In addition, vacuolization and increments in the distance of the intercellular area were observed in the hepatocytes of the same sections (Figure 4).

A severe visceral oedema was detected in larvae at 25 µg L<sup>-1</sup> in the cross-sections of the internal organs (Figure 5). As only one larva survived at 50 µg L<sup>-1</sup>, the histopathological findings are not presented here.

The incidence of observed histopathological findings of the ten larvae is presented in Table 4.

## DISCUSSION

There is a strong relationship between increasing concentrations of cadmium and the decline in the hatching success of frog embryos (37). Cadmium LC<sub>50</sub> values determined in previous acute toxicity studies on amphibian larvae vary considerably. The 96-h LC<sub>50</sub> of cadmium for *Bufo arenarum* larvae was reported in the range from 2.19 to 6.77 mg L<sup>-1</sup> (38). The LC<sub>50</sub> values for cadmium in *R. ridibunda* larvae calculated in two different studies were 0.45 mg L<sup>-1</sup> (39) and 71.8 mg L<sup>-1</sup> (40). This last value is very close to the 90-h LC<sub>50</sub> reported in *Xenopus laevis* larvae - 80 to

**Table 1** Lethal concentrations after cadmium exposure in *P. viridis* for 96 hours

Lethal concentrations	Concentration (µg L <sup>-1</sup> )	95 % Confidence interval	
		Lower bound	Upper bound
LC <sub>10</sub>	26.98	21.05	31.24
LC <sub>50</sub>	35.35	30.41	41.09
LC <sub>90</sub>	46.31	40.05	59.36


**Table 2** Comparison of morphological measurements of treatment groups

Measurements	Cd ( $\mu\text{g L}^{-1}$ )	n	mean	SD	SE	95 % Confidence interval for mean		Min.	Max.	p
						Lower bound	Upper bound			
Total length (mm)	0	10	15.76	0.89	0.28	15.12	16.40	13.95	16.77	0.00
	1	10	15.08	1.18	0.37	14.23	15.92	12.72	16.99	
	5	10	14.88	0.75	0.26	14.25	15.51	13.95	15.88	
	10	10	14.78	1.30	0.45	13.69	15.86	13.18	16.47	
	25	10	13.45	1.38	0.48	12.29	14.61	10.70	14.91	
Width (mm)	0	10	3.08	0.31	0.09	2.85	3.30	2.51	3.61	0.69
	1	10	3.20	0.85	0.27	2.59	3.82	0.92	4.04	
	5	10	3.03	0.37	0.13	2.72	3.35	2.14	3.27	
	10	10	3.29	0.25	0.09	3.07	3.50	3.04	3.77	
	25	10	2.97	0.36	0.12	2.66	3.27	2.15	3.37	
Tail length (mm)	0	10	9.62	0.78	0.24	9.07	10.18	7.90	10.38	0.34
	1	10	9.19	0.78	0.24	8.64	9.75	7.72	10.11	
	5	10	9.39	0.87	0.35	7.75	9.53	7.54	10.78	
	10	10	9.43	0.87	0.30	8.70	10.15	7.90	10.70	
	25	10	8.78	1.19	0.42	7.79	9.78	6.25	10.20	
Wet weight (g)	0	10	0.03	0.01	0.01	0.02	0.03	0.03	0.04	0.33
	1	10	0.03	0.01	0.01	0.02	0.03	0.03	0.04	
	5	10	0.03	0.01	0.01	0.02	0.03	0.03	0.04	
	10	10	0.03	0.01	0.01	0.02	0.03	0.02	0.04	
	25	10	0.02	0.01	0.01	0.01	0.03	0.01	0.04	

100 mg L<sup>-1</sup> (41), but both were substantially above the LC<sub>50</sub> calculated for *X. laevis* embryos - 850  $\mu\text{g L}^{-1}$ . The cadmium 96-h LC<sub>50</sub> for stage 26 *P. variabilis* larvae in our study was 35.35  $\mu\text{g L}^{-1}$ . The dissimilar lethal concentration values calculated in all of these studies might have been due to differences in environmental conditions in habitats, such as temperature and pH, or perhaps due to general differences in species. The developmental stage at the moment of exposure also influences lethal concentration estimations. In a study on juvenile *R. ridibunda*, the LC<sub>50</sub> calculated after a 96-h exposure to cadmium was 51.2 mg L<sup>-1</sup> (42), a value greater than or similar to the LC<sub>50</sub> found for the larvae of this species. It is therefore important to analyse the water at several stages; not only once. According to Turkish regulations, the highest acceptable value is 100  $\mu\text{g L}^{-1}$  (43), while this value was set at 200  $\mu\text{g L}^{-1}$  in the 2008 TSE limits (the Turkish Standards Institution) (43).

This study detected serious declines in survival percentages of the larvae resulting from cadmium exposure at between 25 and 50  $\mu\text{g L}^{-1}$ . We could say that cadmium levels within 25-50  $\mu\text{g L}^{-1}$  were critical for *P. variabilis* larvae. This is most strongly confirmed by the survival of one larva in the group exposed to 50  $\mu\text{g L}^{-1}$  of cadmium. James and Little (13) reported that 540  $\mu\text{g L}^{-1}$  of chronically applied cadmium

reduced survival and extended the metamorphosis period in *Bufo americanus* larvae, but prematurely induced metamorphosis at 5 and 54  $\mu\text{g L}^{-1}$ . Cadmium exposure at 0.25-5  $\mu\text{g L}^{-1}$  stimulated growth and metamorphosis in *Rana pipiens* larvae (16, 32). In another study (15), chronic exposure to 0-855  $\mu\text{g L}^{-1}$  of cadmium did not affect the survival rate of *X. laevis* larvae. When discussing the complexity of amphibian larval development (under extensive hormonal regulation), it may be concluded that small concentrations of cadmium exposure may even have a stimulating effect on this process.




**Figure 2** Cross sections of the gill lamellae of (a) control group, (b) 10  $\mu\text{g L}^{-1}$  cadmium, and (c) 25  $\mu\text{g L}^{-1}$  cadmium (lamellar fusion shown with arrows), H&E staining



**Table 3** Statistically significant values of the discriminant analysis made so as to reveal the difference among the groups following cadmium exposure in *P. viridis* larvae for 96 hours

Function	Eigenvalue	Variance (%)	Total (%)	Canonical Correlation	Wilks Lambda	Chi-square	df	p
1	0.648	94	94	0.627	0.583	24.556	8	0.00
2	0.041	6	100.0	0.199	0.960	1.835	3	0.60




**Figure 3** Cross sections of the pronephric tubules of (a) control group, and (b) 25  $\mu\text{g L}^{-1}$  cadmium (deformations of pronephric tubules shown with asterisk), H&E staining


**Table 4** The incidence of observed histopathological findings in the gill, kidney, and liver of larvae administered cadmium

Lesions	Cadmium ( $\mu\text{g L}^{-1}$ )				
	0	1	5	10	25
Gills lamellar fusions	0	0	0	5	7
Kidney deformation of pronephric tubules	0	0	1	3	6
Liver deformation and haemorrhage	0	0	1	3	6
Vacuolisation	0	0	0	6	8
Visceral oedema	0	0	3	5	8

$n=10$ . Values indicate the numbers of larvae with observed lesions in their sections



**Figure 4** Cross sections of the liver of (a) control group, (b) 10  $\mu\text{g L}^{-1}$  cadmium and (c) 25  $\mu\text{g L}^{-1}$  cadmium (haemorrhage shown with asteriks, vacuolization shown with arrows), H&E staining



**Figure 5** Cross sections of the internal organs of (a) control group, and (b) 25  $\mu\text{g L}^{-1}$  cadmium (visceral oedema shown with asterisk), H&E staining

Heavy metal exposure at sub-lethal concentrations initially affects the gills, with the potential risk of accumulation (44). Gill damage, in particular fusion of secondary lamellae, has been reported in *Channa punctatus* exposed to 172  $\text{mg L}^{-1}$  of cadmium during 10 days (45). The most important cause of gill lamellar fusion due to pollutant exposure is their function as a barrier for preventing pollutant entry into the body (46). Likewise, fusions in gill lamellae in our study were found in larvae exposed to 10 and 25  $\mu\text{g L}^{-1}$  of cadmium.

Cadmium causes histopathological changes in the liver, kidney, gill, spleen, and bone marrow, as well as hypocalcemia and hypoglycemia, and inhibits the intake of  $\text{Ca}^{2+}$  through gills, therefore affecting plasma ion composition and osmoregulation (27). It has also been reported that cadmium has hepatotoxic effects and causes hepatic dysfunctions and increases in certain enzyme activities (14), as well as histopathological lesions in the liver and necrosis in hepatocytes (45, 47). Furthermore, a study on *Bufo arenarum* revealed that cadmium accumulates particularly in the liver (48). Much like in other sources, our study also detected serious deformations in the hepatic and renal tubules. The damage observed in liver cross-sections, necrosis, lesions, and haemorrhages verify that cadmium primarily targets the liver. Effects on kidneys include epithelial cell deformations in the pronephric tubules and enlargement of the tubular lumen. Similar results were reported for *Rana ridibunda* in a study that revealed progressive nephropathy and glomerulonephropathy as a result of cadmium exposure (14).

In conclusion, even though the histopathological findings of our study support the results of previous studies, its main contribution is that it is the first to report results for *P. variabilis*, which has shown to be a very good indicator species for aquatic ecosystems. Our  $\text{LC}_{50}$  values were lower than those reported by

other investigators. The mechanisms and biological route of cadmium, with special reference to amphibians, has not yet been fully clarified and further detailed acute, sublethal, and chronic toxicity studies are necessary.

## REFERENCES

1. Halliday T. Diverse phenomena influencing amphibian population declines. In: Lannoo MJ, editor. Amphibian declines - The conservation status of United State species. Berkeley (CA): University of California Press; 2005. p. 3-6.
2. Barnum TR, Verburg P, Kilham SS, Whiles MR, Lips KR, Colon-Gaud C, Pringle CM. Use of stable isotope ratios to characterize potential shifts in the isotopic niches of grazing insects following an amphibian decline in a Neotropical stream. *J Trop Ecol* 2013;29:291-9. doi: 10.1017/S026646741300028X
3. Collins JP, Storfer A. Global amphibian declines: sorting the hypotheses. *Divers Distrib* 2003;9:89-98. doi: 10.1046/j.1472-4642.2003.00012.x
4. Rohr JS, Raffel TR. Linking global climate and temperature variability to widespread amphibian declines putatively caused by disease. *Proc Natl Acad Sci USA* 2010;107:8269-74. doi: 10.1073/pnas.0912883107
5. Clark EJ, Norris DO, Jones RE. Interactions of gonadal steroids and pesticides (DDT, DEE) on gonaduct growth in larval tiger salamanders, *Ambystoma tigrinum*. *Gen Comp Endocrinol* 1998;109:94-105.
6. Goleman W, Urguidi LJ, Anderson TA, Smith EE, Kendall RJ, Carr JA. Environmentally relevant concentrations of ammonium perchlorate inhibit development and metamorphosis in *Xenopus laevis*. *Environ Toxicol Chem* 2002;21:424-30. doi: 10.1002/etc.5620210227
7. Qin ZF, Zho JM, Chu SG, Xu XB. Effects of Chinese domestic polychlorinated biphenyls (PCBs) on gonadal differentiation in *Xenopus laevis*. *Environ Health Perspect* 2003;111:553-6. doi: 10.1289/ehp.5620
8. Bernabo I, Brunelli E, Berg C, Bonacci A, Tripepi S. Endosulfan acute toxicity in *Bufo bufo* gills: ultrastructural changes and nitric oxide synthase localization. *Aquat Toxicol* 2008;86:447-56. doi: 10.1016/j.aquatox.2007.12.006

9. Cooke AS. Tadpoles as indicators of harmful levels of pollution in the field. *Environ Pollut* 1981;25:123-33. doi: 10.1016/0143-1471(81)90012-X
10. Lajmanovich RC, Sanches-Hernandes JC, Peltzer PM, Attademo AM, Fiorenza MC, Cabagna MC, Basso A. Levels of plasma B-esterase and Glutathione-S-transferase activities in three South American toads. *J Toxicol Environ Chem* 2008;90:1145-61. doi: 10.1080/02772240801923107
11. Zhelev ZM, Popgeorgiev GS, Angelov MV. Investigating the changes in the morphological content of the blood of *Pelophylax ridibundus* (Amphibia: Ranidae) as a result of the anthropogenic pollution and its use as an environmental bioindicator. *Acta Zool Bulg* 2013;65:187-96.
12. Flament S, Kuntz S, Chesnel A, Grillier-Vuissoz I, Tankozic C, Penrad-Mobayed M, Auque G, Shiralı P, Schroeder H, Chardard D. Effects of cadmium on gonadogenesis and metamorphosis in *Pleurodeles waltl* (urodele amphibian). *Aquat Toxicol* 2003;64:143-53. doi: 10.1016/S0166-445X(03)00042-0
13. James SM, Little EE. The effects of chronic cadmium exposure on American toad (*Bufo americanus*) tadpoles. *Environ Toxicol Chem* 2003;22:377-80. doi: 10.1002/etc.5620220219
14. Loumbourdis NS. Hepatotoxic and nephrotoxic effects of cadmium in the frog *Rana ridibunda*. *Arch Toxicol* 2005;79:434-40. doi: 10.1007/s00204-005-0652-x
15. Sharma B, Patino R. Exposure of *Xenopus laevis* tadpoles to cadmium reveals concentration-dependent bimodal effects on growth and monotonic effects on development and thyroid gland activity. *Toxicol Sci* 2008;105:51-8. doi: 10.1093/toxsci/kfn119
16. Gross JA, Johnson PTJ, Prahı LK, Karasov WH. Critical period of sensitivity for effects of cadmium on frog growth and development. *Environ Toxicol Chem* 2009;28:1227-32. doi: 10.1897/08-205.1
17. Jolibois LS, Shi LS, George W, Henson WJ, Anderson MB. Cadmium accumulation and effects on progesterone release by cultured human trophoblast cells. *Reprod Toxicol* 1999;13:215-21. doi: 10.1016/S0890-6238(99)00009-X
18. McCarthy JF, Shugart LR, editors. Biomarkers of Environmental Contamination. Boca Raton (FL): Lewis Publishers; 1990.
19. Satarug S, Baker JR, Urbenjapol S, Haswell-Elkins M, Reilly PEB, Williams DJ, Moore MR. A global perspective on cadmium pollution and toxicity in non-occupationally exposed population. *Toxicol Lett* 2003;137:65-83. doi: 10.1016/S0378-4274(02)00381-8
20. Altındağ A, Yiğit S. Assessment of heavy metal concentrations in the food web of lake Beyşehir, Turkey. *Chemosphere* 2005;60:552-6. doi: 10.1016/j.chemosphere.2005.01.009
21. Baker JR, Satarug S, Urbenjapol S, Edwards RJ, Williams DJ, Moore MR, Reilly PEB. Potential for early involvement of CYP isoforms in aspects of human cadmium toxicity. *Toxicol Lett* 2003;137: 85-93. doi: 10.1016/S0378-4274(02)00382-X
22. Jarub L, Akesson A. Current status of cadmium as an environmental health problem. *Toxicol Appl Pharmacol* 2009;238: 201-8. doi: 10.1016/j.taap.2009.04.020
23. Ranatunge RAAR, Wijesinghe MR, Ratnasooriya WD, Dharmarathne HASG, Wijesekera RD. Cadmium-induced toxicity on larvae of the common asian toad *Duttaphrynus melanostictus* (Schneider 1799): Evidence from emperial trials. *Bull Environ Contam Toxicol* 2012;89:143-6. doi: 10.1007/s00128-012-0635-6
24. Weltje L, Simpson P, Gross M, Crane M, Wheeler JR. Comparative acute and chronic sensitivity of fish and amphibians: A critical review of data. *Environ Toxicol Chem* 2013;32:984-94. doi: 10.1002/etc.2149
25. Gay F, Laforgia V, Caputo I, Esposito C, Lepretti M, Capaldo A. Chronic exposure to cadmium disrupts the adrenal gland activity of the newt *Triturus carnifex* (Amphibia, Urodela). *BioMed Res Int* 2013;1-6. doi: 10.1155/2013/424358
26. Spry DJ, Wiener JG. Metal bioavailability and toxicity to fish in low-alkalinity lakes; A critical review. *Environ Pollut* 1991;71:243-304. doi: 10.1016/0269-7491(91)90034-T
27. Ricard AC, Daniel C, Anderson P, Hontela A. Effects of subchronic exposure to cadmium chloride on endocrine and metabolic functions in rainbow trout *Oncorhynchus mykiss*. *Arch Environ Contam Toxicol* 1998;34:377-81. doi: 10.1007/s002449900333
28. Vetillard A, Bailhache T. Cadmium: An endocrine disrupter that affects gene expression in the liver and brain of juvenile rainbow trouts. *Biol Reprod* 2005;72:119-26. PMID: 15317685
29. Calevro F, Campani S, Ragghianti M, Bucci S, Mancino G. Tests of toxicity and teratogenicity in biphasic vertebrates treated with heavy metals (Cr<sup>3+</sup>, Al<sup>3+</sup>, Cd<sup>2+</sup>). *Chemosphere* 1998;37:3011-7. doi: 10.1016/S0045-6535(98)00342-7
30. Mouchet F, Baudrimont M, Gonzalez P, Cuenot Y, Bourdineaud JP, Boudou A, Gauthier L. Genotoxic and stress inductive potential of cadmium in *Xenopus laevis* larvae. *Aquat Toxicol* 2006;78:157-66. doi: 10.1016/j.aquatox.2006.02.029
31. Zhang Y, Huang D, Zhao D, Long J, Song G, Li A. Long-term toxicity effects of cadmium and lead on *Bufo raddei* tadpoles. *Bull Environ Contam Toxicol* 2007;79:178-83. doi: 10.1007/s00128-007-9152-4
32. Gross JA, Chen TH, Karasov WH. Lethal and sublethal effects of chronic cadmium exposure on northern leopard frog (*Rana pipiens*) tadpoles. *Environ Toxicol Chem* 2007;26:1192-7. doi: 10.1897/06-479R.1
33. Başoğlu M, Özeti N, Yılmaz I. Türkiye Amfibileri [The Amphibians of Turkey, in Turkish]. İzmir: Ege Üniversitesi Fen Fakültesi; 1992.
34. Kinzelbach R, Kasperek M. Amphibia and Reptilia. *Zool Middle East* 1992;3:27.
35. International Union for Conservation of Nature (IUCN). The IUCN Red List of Threatened Species. [Displayed 25 August 2014]. Available at <http://www.iucnredlist.org/>
36. Gosner KL. A simplified table for staging anuran embryos and larvae with notes on identification. *Herpetologica* 1960;16:183-90.
37. Haywood LK, Alexander GJ, Byrne MJ, Cukrowska E. *Xenopus laevis* embryos and tadpoles as models for testing pollution by zinc, copper, lead and cadmium. *Afr Zool* 2004;39:163-74.
38. Ferrari L, Saliban A, Muino CV. Selective protection of temperature against cadmium acute toxicity of *Bufo arenarum* tadpoles. *Bull Environ Contam Toxicol* 1993;50:212-8. doi: 10.1007/BF00191724
39. Grillitch B, Chovanec A. Heavy metals and pesticides in anuran spawn and tadpoles, water, and sediment. *Toxicol Environ Chem* 1995;50:131-55. doi: 10.1080/02772249509358210



40. Loumbourdis NS, Kyriakopoulou-Sklavounou P, Zachariadis G. Effects of cadmium exposure on bioaccumulation and larval growth in the frog *Rana ridibunda*. Environ Pollut 1999;104:429-33. doi: 10.1016/S0269-7491(98)00172-9
41. Woodal C, Maclean N, Crossley F. Responses of the trout fry (*Salmo gairdneri*) and *Xenopus laevis* tadpoles to cadmium and zinc. Comp Biochem Physiol C 1988;89:93-9. doi: 10.1016/0742-8413(88)90151-X
42. Selvi M, Gül A, Yılmaz M. Investigation of acute toxicity of cadmium chloride ( $\text{CdCl}_2 \cdot \text{H}_2\text{O}$ ) metal salt and behavioral changes it causes on water frog (*Rana ridibunda* Pallas, 1771). Chemosphere 2003;52:259-63. doi: 10.1016/S0045-6535(03)00262-5
43. Kayhan FE. Su ürünlerinde kadmiyum biyobirikimi ve toksisitesi [Bioaccumulation and toxicity of cadmium in the water products, in Turkish]. J Fish Aquat Sci 2006;23:215-20.
44. Flos R, Tort L, Balasch J. Effects of zinc sulphate on haematological parameters in the dogfish *Scyliorhinus canicula* and influences of MS222. Mar Environ Res 1987;21:289-98. doi: 10.1016/0141-1136(87)90051-1
45. Babu RA. Effects of cadmium on digestive organs of teleost fish *Ophiocephalus* (Channa). Universal J Environ Res Technol 2013;3:173-80.
46. Mallat J. Fish gill structural changes induced by toxicants and other irritants: A statistical review. Can J Fish Aquat Sci 1985;42:630-48. doi: 10.1139/f85-083
47. Mitsumori K, Shibutani M, Sato S, Onodera H, Nakagawa J, Hayashi Y, Ando M. Relationship between the development of hepato-renal toxicity and cadmium accumulation in rats given minimum to large amounts of cadmium chloride in the long-term: Preliminary study. Arch Toxicol 1998;72:545-52. doi: 10.1007/s002040050541
48. Pérez-Coll CS, Herkovits J, Salibian A. [Effects of cadmium on the development of an amphibian, in Spanish]. Arch Biol Med Exp (Santiago) 1985;18:33-9. PMID: 3939390



### Sažetak

#### Akutni toksični učinci kadmija na ličinke zelene žabe, *Pseudepidalea variabilis* (Pallas, 1769) (Amphibia: Anura)

Štetni utjecaji kadmija na okoliš i njegovo nakupljanje u prirodi posljednjih su godina poprimili zabrinjavajuće razmjere. U okviru ove studije istražili smo akutne toksične učinke kadmija na morfologiju i histologiju ličinaka zelene žabe, *Pseudepidalea variabilis* (Pallas, 1769). Embriji žabe dobiveni su od jedinki koje su u fazi amplexusa prikupljene u prirodi. U laboratorijskim uvjetima embriji su bili držani do razvojnog stadija 26, kada su izloženi kadmiju u koncentracijama od 0, 1, 5, 10, 25 i 50  $\mu\text{g L}^{-1}$  tijekom 96 sati. U pokusu smo odredili sljedeće letalne koncentracije kadmija:  $\text{LC}_{10}$  26,98  $\mu\text{g L}^{-1}$ ,  $\text{LC}_{50}$  35,35  $\mu\text{g L}^{-1}$  i  $\text{LC}_{90}$  46,31  $\mu\text{g L}^{-1}$ . Izloženost kadmiju u koncentracijama većim od 1  $\mu\text{g L}^{-1}$  negativno je utjecala na veličinu tijela ličinaka. Histološke analize upućuju na sljepljivanje škržnih listića, krvarenja u jetrima, pojavu edema u trbušnoj šupljini i deformacije pronefričkih kanalića (pri koncentracijama većim od 10  $\mu\text{g L}^{-1}$ ). Dobiveni rezultati pokazali su da je zelena žaba vrlo osjetljiva na kadmij, na što upućuje vrijednost  $\text{LC}_{50}$  koja je u našem pokusu bila niža od vrijednosti zabilježenih u drugim istraživanjima. Prema tome, ta se vrsta može smatrati pouzdanim pokazateljem okolišnog stresa u slatkovodnim ekosustavima.

**KLJUČNE RIJEČI:** embriotoksičnost; histopatologija; letalne koncentracije; teški metal

#### CORRESPONDING AUTHOR:

Sibel Hayretdağ  
Terzioğlu Campus,  
17100 Çanakkale, Turkey  
E-mail: [sibelhayretdag@gmail.com](mailto:sibelhayretdag@gmail.com)